ERRATA for Diagnostic Radiology Coding Reference 2015 Edition

Text deletions are crossed out. New text is **blue and bolded**. Ordered by appearance in text.

Page 65, Coding Instructions

11. The 2015 *CPT Codebook* recommends use of code 77003 for fluoroscopic imaging guidance and localization with code 62284 or 61055 when a formal contrast radiographic myclogram study is not performed. Code 77003 is bundled with all imaging codes that utilize contrast (70000 series) codes 72240-72270 and as well as codes 62302-62305.

Page 103, Coding Instructions

2. When a screening mammogram is performed that shows an abnormality requiring diagnostic films, and the additional films are taken while the patient is still there, report both codes for the screening mammogram and the diagnostic mammogram. Append modifier -GG to the diagnostic mammogram to indicate both were done on the same day due to findings on the screening study. At this time a -59 modifier must also be appended to the screening mammography code to satisfy an NCCI edit that modifier -GG will not pass. In this scenario, the radiologist can order the diagnostic mammogram. If the patient has already left the facility when it is determined that a diagnostic mammogram is needed, a new order from the patient's treating physician for the diagnostic mammogram must be obtained.

Page 106, Coding Instructions

1. When a screening mammogram is performed that shows an abnormality requiring diagnostic films, and the additional films are taken while the patient is still there, report both codes for the screening mammogram and the diagnostic mammogram. Append modifier -GG to the diagnostic mammogram to indicate both were done on the same day due to findings on the screening study. At this time a -59 modifier must also be appended to the screening mammography code to satisfy an NCCI edit that modifier -GG will not pass. In this scenario, the radiologist can order the diagnostic mammogram. If the patient has already left the facility when it is determined that a diagnostic mammogram is needed, a new order from the patient's treating physician for the diagnostic mammogram must be obtained.